

**THE EFFICACY OF AEROBIC EXERCISE AND RELAXATION
TECHNIQUE IN FATIGUE AMONG CHILDREN WITH ACUTE
LYMPHOBLASTIC LEUKEMIA- A COMPARATIVE STUDY**

Dissertation submitted to

The Tamil Nadu Dr. M.G.R. Medical University

Chennai

In partial fulfillment of the requirements for the degree of

MASTER OF PHYSIOTHERAPY

(PEDIATRICS)



Reg. No. 271540062

OCTOBER – 2017

**COLLEGE OF PHYSIOTHERAPY
SRI RAMAKRISHNA INSTITUTE OF PARAMEDICAL SCIENCES
COIMBATORE – 641044**

CERTIFICATE

This is to certify that the dissertation work entitled **“The Efficacy of Aerobic Exercise and Relaxation Technique in Fatigue Among Children with Acute Lymphoblastic Leukemia- A Comparative Study”** was carried out by the candidate bearing the **Register No.271540062 (October 2017)** in College of Physiotherapy, SRIPMS, Coimbatore, affiliated to the Tamil Nadu Dr. M.G.R Medical University, Chennai towards partial fulfillment of the **Master of Physiotherapy (Pediatrics)**.

Prof. B. SANKAR MANI, MPT (Sports),MBA,
Principal
College of Physiotherapy
SRIPMS
Coimbatore – 641044

Place: Coimbatore

Date:

CERTIFICATE

This is to certify that the dissertation work entitled **“The Efficacy of Aerobic Exercise and Relaxation Technique in Fatigue Among Children with Acute Lymphoblastic Leukemia- A Comparative Study”** was carried out by the candidate bearing the **Register No.271540062 (October 2017)** in College of Physiotherapy, SRIPMS, Coimbatore, affiliated to the Tamil Nadu Dr. M.G.R Medical University, Chennai towards partial fulfillment of the **Master of Physiotherapy(Pediatrics)** under my direct supervision and guidance.

**Prof .V. SUTHAKAR, MPT (Pediatrics),
Guide**

College of physiotherapy
SRIPMS
Coimbatore – 641044

Place: Coimbatore

Date:

CERTIFICATE

This is to certify that the dissertation work entitled “**The Efficacy of Aerobic Exercise and Relaxation Technique in Fatigue Among Children with Acute Lymphoblastic Leukemia - A Comparative Study**” was carried out by the candidate bearing the **Register No.271540062 (October 2017)** College of Physiotherapy, SRIPMS, Coimbatore affiliated to The Tamilnadu Dr. M.G.R Medical University, Chennai towards partial fulfillment of the requirements for the degree of **Master of Physiotherapy (Paediatrics)** was evaluated.

INTERNAL EXAMINER

EXTERNAL EXAMINER

Place:

Date:

ACKNOWLEDGEMENT

With great privilege I express my deep sense of gratitude to the *God Almighty* for his blessings, love and care for me and who have always been my source of inner strength and courage throughout my life.

From the bottom of my heart, I thank **my dear Parents, my Brother** and family **Members** for their concern and endless love for me in every part of my life.

With great honor, I dedicate this study to my guide **Prof.V.Suthakar, MPT (Pediatrics)**, College of Physiotherapy, SRIPMS, Coimbatore, without whom this venture would have been impracticable. I am grateful to his inevitable role in organizing and completing the study with full support and patience.

My sincere thanks to **Professor .Mr. B. Sankarmani, MPT (Sports), MBA., Principal**, College of Physiotherapy, SRIPMS, Coimbatore, who provided me the opportunity to perform the study.

I am grateful to **Dr.Guhan, Chief Oncologist, Sri Rama Krishna Institute of Oncology and Research** for his acceptance in conducting this study.

I extend my sense of gratitude to all **the Staffs** of College of Physiotherapy, SRIPMS, for their timely help and valuable information for the betterment of my study.

I am propitious to have such a great bunch of **Parents, Caregivers** and **staff nurses** who made me trustworthy to involve their precisions and no words could be enough to convey my boundless love to each of the **little Ones** who made the study colorful.

My sincere gratification to all the **non-teaching staffs** of Sri Ramakrishna Institute who have been a part of this study.

A token of appreciation to my friends **Anjana, Surya, Bhavitra** and **my classmates** for their helping hands at the right time.

I thank **Saraswathi Computer Centre**, Coimbatore, in particular for their defined and orderly execution of the dissertation work.

ABBREVIATIONS

ALL	-	Acute Lymphoblastic Leukemia
CRF	-	Cancer Related Fatigue
CFS	-	Child Fatigue Scale
PMR	-	Progressive Muscle Relaxation
PFS	-	Parent Fatigue Scale
QOL	-	Quality Of Life

CONTENTS

No.	Title	Page no.
1.	Introduction	1
2.	Objectives	7
3.	Review of Literature	10
4.	Methodology	14
5.	Data analysis & Results	20
6.	Discussion	30
7.	Conclusion	34
	References	35
	Annexures	

LIST OF TABLES

No.	Title	Page no.
I.	CFS scoring for Group A and B	22
II.	PFS scoring for Group A and B	24
III.	CFS scoring for Group A vs. B	26
IV.	PFS scoring for Group A vs. B	28

LIST OF GRAPHS

No.	Title	Page no.
I.	CFS scoring for Group A and Group B	22
II.	PFS scoring for Group A and Group B	24
III.	CFS scoring in Group A vs. Group B	26
IV.	PFS scoring in Group A vs. Group B	28

1. INTRODUCTION

Childhood cancer is one of the leading cause of death by disease among which Acute Lymphoblastic Leukemia (ALL) accounts for 30% of all the malignancies⁽¹⁾. ALL is the most common form of pediatric leukemia which is also known as Acute Lymphocytic or Acute Lymphoid Leukemia⁽²⁾. Incidence is approximately 3-4 cases per 100,000 children below 15 years with peak incidence below 2-5 years of age⁽³⁾. In most of the countries, the incidence rate in children was approximately four times than that of the adults⁽⁴⁾. Studies have shown a 98% cure rate in ALL⁽⁵⁾. But even with these survival rates in pediatric population, only half of the ALL patients survived 5 years⁽⁶⁾.

Exact etiology of ALL is still unknown but common etiological factors include ionizing radiations ,congenital anomalies, chemical agents like Benzene and Immunodeficiency syndromes⁽⁷⁾. Many studies have reported the risk factors like Down syndrome, Erbstien Barr Virus (EBV) infections, parental history of fetal loss, miscarriage, perinatal exposure to pesticides and still birth to be reasons for developing leukemia^(8, 9, 10).

The pathogenesis of ALL is assumed to be the neoplasm arising from the clonal proliferation of immature cells of hemopoetic system which are characterized by the aberrant or arrested differentiation ⁽⁷⁾.

The associated complications of ALL include musculoskeletal, Central Nervous System (CNS) and other systemic complications. Recent study have shown that 2% of the boys diagnosed with ALL shows testicular enlargement ⁽⁵⁾. Neural complications like pain, paresthesia and reduced deep tendon reflexes have been reported in various studies ⁽¹¹⁾. Musculoskeletal complications like impaired gross motor and fine motor performances, increased fatigue, compromised ROM, strength and endurance have also been identified^(11,12).

Unfortunately, fatigue has become the most debilitating symptom in patients with cancer ⁽¹³⁾. It was reported to be an intensive disability among cancer patients ⁽¹⁴⁾ as it induces inactivity which on prolonged rest lead to muscle wasting and loss of endurance ⁽¹²⁾. Fatigue affects quality of life to about 60% in cancer patients. But limited focus was given to those groups recognized with fatigue complaining unique experiences ⁽¹⁵⁾.

The medical management of ALL includes antineoplastic agents administered for induction, consolidation and maintenance therapy ⁽¹⁶⁾. For newly diagnosed form of ALL, Asparagine and Anthracycline are administered ⁽⁵⁾. Optimal use of antileukemic agents together with prognostic factors resulted in steady improvement of treatment outcome ⁽¹⁷⁾. Findings have shown that even after excluding cranial irradiation approximately 6% of children with ALL may relapse after completion of treatment and those who remain in remission at 4 years is considered cured ⁽³⁸⁾. Since most of the deaths were assumed to be infection related, corticosteroids, prophylactic antibiotics and antifungals are administered for preventing infections in children receiving chemotherapy ^(16, 18).

Many studies have been undergone and is still on work regarding the hazardous side effects of drugs on ALL. Studies have reported the lower doses of Asparagine for longer periods were responsible for high incidence of thrombosis ⁽¹⁹⁾. Dexamethasone effect on treating pediatric ALL was found effective but caused serious metabolic and neurophysiological side effects ⁽²⁰⁾. Various studies focused on greater risk of developing Osteonecrosis and myocardial impairment in children receiving Corticosteroids and

Doxorubicin treatment ^(21, 22). Dosage of methotrexate was found to cause neurocognitive impairment ⁽²³⁾. Associated mobility functions and weakness were exhibited by children undergoing treatment at standard risk ⁽²⁴⁾. The age of impact was studied to be one of the most important prognostic factor for outcome ⁽²⁵⁾. Thus an early diagnosis, infection control and parental education is needed to improve the conditions of children and many studies have pointed the importance of a feasible intra hospital or home based rehabilitation program for cancer children⁽²⁶⁾.

The rehabilitation services for musculoskeletal, neurological and cardiovascular late effects among ALL children proved to improve the physical function ⁽²⁷⁾. But for an effective physical therapy intervention age, type of exercises, location of implementation and parent motivations are necessary ⁽²⁸⁾. Many studies clarified the extend of physical therapy to be implemented should be within the normal safe boundaries ⁽²⁴⁾. There are studies providing strong evidences of implementing exercises which brought noticeable effect even during chemotherapy^(29,30). These exercises showed appreciable improvement in level of fatigue and endurance improving Quality of Life (QOL)^(2, 31).

Evaluation of presence of fatigue should be clearly made prior to treatment through regular screening and assessment ⁽³²⁾. Numerous studies have been carried out to identify a valid scale to measure fatigue level in children and was put forward as reliable enough^(32,33,34). Many reports proved the effectiveness of aerobic exercise in children with ALL and even during the chemotherapy ^(14, 30, 31). These training focused to improve strength, functional mobility, reduce fatigue, and improve cardiovascular fitness, improve muscle strength and physical function ^(35, 36). Relaxation techniques implemented on children brought greater results than aerobic exercise in certain studies thereby reducing fatigue level and cancer related symptoms ⁽³⁷⁾.

NEED FOR THE STUDY

There are various physiotherapy treatments available for management of ALL related fatigue thereby improving endurance and QOL .In spite of all these managements, the improvement in the level of fatigue was not satisfactory. Fatigue measures for children and adolescents are important because they allow degree of fatigue to be described and high risk groups to be identified ⁽³⁹⁾.

There are studies reporting the relationship between aerobic exercise, relaxation technique and fatigue which showed greater results. Some studies also showed that exercise was not as effective as expected to bring down the fatigue level. On the other hand strong evidences supporting the exercise interventions on relieving fatigue level have been stated without much adverse effects .This shows that the effect of relaxation technique and aerobic training have to be studied further for clinical evidence.

AIM OF THE STUDY

This study was aimed to find out the effect of aerobic training and relaxation technique on fatigue in children with ALL.

2. OBJECTIVES

- To find out the effectiveness of CSF in aerobic training among children with ALL.
- To find out the effectiveness of PSF in aerobic training among children with ALL.
- To find out the effectiveness of CSF in relaxation technique among children with ALL.
- To find out the effectiveness of PSF in relaxation technique among children with ALL.
- To compare the outcomes of CSF between the aerobic training and relaxation technique among children with ALL.
- To compare the outcomes of PSF between the aerobic training and relaxation technique among children with ALL.

HYPOTHESIS

Null Hypothesis

1. There was no significant difference in CFS outcome in aerobic training among children with ALL.
2. There was no significant difference in PFS outcome in aerobic training among children with ALL.

3. There was no significant difference in CFS outcome in relaxation technique among children with ALL.
4. There was no significant difference in PFS outcome in relaxation technique among children with ALL.
5. There was no significant difference in outcome of CFS between the aerobic training and relaxation technique among children with ALL.
6. There was no significant difference in outcome of PFS between the aerobic training and relaxation technique among children with ALL.

Alternate Hypothesis

1. There was a significant difference in CFS outcome in aerobic training among children with ALL.
2. There was a significant difference in PFS outcome in aerobic training among children with ALL.
3. There was a significant difference in CFS outcome in relaxation technique among children with ALL.
4. There was significant difference in PFS outcome in relaxation technique among children with ALL.

5. There was a significant difference in outcome of CFS between aerobic training and relaxation technique group.
6. There was a significant difference in outcome of PFS between aerobic training and relaxation technique group.

.

3. REVIEW OF LITERATURE

Dhoriyani Narendra B, *et al*⁽⁷⁾ concluded that relaxation exercises proved to be more effective in treating fatigue among ALL children than aerobic exercise as it relieves stress and was easy to perform among groups of children that lasted for 3 weeks.

Fernando. C. Dimeo, *et al*⁽³⁷⁾ concluded the effects of aerobic training which could improve the physical performance but was no better than progressive relaxation technique in treating fatigue.

Ali Hassan pour Dekhordi and AmisJalali⁽⁴⁶⁾ in their study pointed out the importance of PMR that had greater effect in relieving fatigue and improving the quality of life.

Tseng Tien Haung and Kirsten K Ness⁽³⁶⁾ summarized the impact of exercise on health and physical function among children during and after cancer treatment. They reported reduced levels of fatigue and improved physical activity in children who completed aerobic exercise program.

Dimeo, *et al*⁽⁴¹⁾ concluded the effects of aerobic exercise program which can be prescribed in defined intensity, duration, frequency as a therapy in managing fatigue among cancer patients.

Shadi Farzin Gohar MD, et al ⁽²⁶⁾ demonstrated the feasibility of physical therapy program in hospital and home based plan during initial four phases of medical treatment and focused on the importance of further studies on the same for better results.

Anna Spathis, et al ⁽¹⁵⁾ concluded fatigue as a major disabling factor among cancer patients in their study and suggested to focus on the effectiveness of interventions to promote activity and manage symptoms

Deborah Tomlison et al ⁽³⁹⁾ in their study analyzed the psychometric properties of the scales measuring fatigue and found the parent, child fatigue scale to be effective.

Dimeo F, et al ⁽¹²⁾ also concluded that aerobic training can be safely carried after high dose chemotherapy thereby preventing loss of physical performance.

Marja Schoenmakers et al ⁽²⁴⁾ conducted the pilot study to find the effects of exercises on muscle strength before and after treatment of ALL and concluded that certain muscle groups showed improvement in activity.

Marilyn .J. Hockenberry, et al⁽³³⁾ in his study to develop and test the instruments for measuring fatigue thereby studying the instrument development and content validation was the first to provide PFS,CFS,SFS as valid and reliable instruments to measure fatigue in children.

Hinds P S, et al⁽³⁹⁾ determined the effects of exposure to dexamethasone on sleep and fatigue in pediatric patients with ALL showing the validity of FSC (Fatigue Scale Children) and FSP (Fatigue Scale Parent).

Donald. P. Lawrence, et al⁽⁴⁸⁾ has concluded that the assessment properties of the scale was only promising and pointed to focus on the development of much precise assessment tools.

Alison Crichton, et a⁽⁴⁷⁾ has appreciated the robust measurement properties notably content validity showed by FSC and FSP scoring systems, providing moderate evidence for structural validity of child and parent forms. The evidence assessed was exclusively limited for children with hematology /oncology diagnosis.

Victoria G Marghese et al⁽¹¹⁾ focused on early initiation of treatment with greater emphasis on endurance activities which improves the quality of life.

San Juan *et al* ⁽³⁵⁾ stated the effects of an intra-hospital exercise program which brought significant increase in measure of aerobic fitness, strength and functional mobility.

Ladha *et al* ⁽⁴⁰⁾ showed that a 30 minute of moderate intensity exercise in ALL children receiving medical treatment provides a similar neutrophil response that of healthy age group.

4. METHODOLOGY

Study design

A comparative study

Study setting

Pediatric inpatient oncology ward, Sri Ramakrishna Institute of Oncology and Research, Coimbatore.

Sampling technique

A convenience sampling.

Sample size

A total of 30 acute lymphoblastic leukemia patients was selected on the basis of inclusion criteria and conveniently allocated into two groups of 15 each. Both the groups were designed for experimentation.

Study duration

The study was conducted over a period of 10 months.

Treatment duration

The treatment duration was 4 weeks.

Inclusion criteria

- Children undergoing chemotherapy
- Aged between 5-12
- Both gender
- Children with preserved cardiac function
- Children able to accept and follow verbal instructions
- Children complaining of fatigue

Exclusion criteria

- Children with unstable cardiac function
- Children with platelet count less than 3000/uL
- Children with recent surgeries(less than 3 months)
- Children with congenital musculoskeletal anomalies
- Children with visual and hearing deficits
- Children with systemic complications.

Method of collection of data

The selected ALL patients were conveniently assigned into two equal groups: Group A/aerobic exercise group and Group B/Relaxation technique group. Both the groups were explained about the study and a written consent (*Annexure I*) was signed and obtained from the parents of children involved in the study.

All the participants were assessed by the pediatric assessment (*Annexure II*) for the general health status which consist of age, gender, height, weight, duration of chemotherapy, vitals and general physical examination.

All the participants were evaluated by the Child Fatigue Scale, (*Annexure III*) which was explained by the trainer to the children and were asked to rate their level of fatigue before and after the completion of the treatment regimen. The parents of all the participants were given with the Parent Fatigue Scale questionnaire, (*Annexure IV*) and was asked to answer it based on the tiredness level of their child before and after the treatment and the total score was taken for analysis.

The group A and B received Aerobic exercise (*Annexure-V*) and Relaxation technique (*Annexure-VI*) respectively for a period of 4weeks. All the vital parameters were measured during each session to confirm that subjects were stable and the treatment sessions were carried out under the presence of the parent.

Outcome Measures

The assessment tools used to assess the level of fatigue were:

1. Child Fatigue Scale
2. Parent Fatigue Scale

TREATMENT TECHNIQUES

Aerobic training group (*Annexure V*)

The children in this group received the low intensity aerobic training which consisted of general warm ups, static cycling, walking on treadmill incorporated with aerobic play activities and finally cool down.

The activities were demonstrated to the children by the therapist. The treatment focused on improving their strength and endurance there by reducing fatigue. The treatment was conveniently arranged for the children which lasts for about 30 minutes a day for 3 alternate days a week for 4 weeks.

The treatment circuit consists of:

- Initial warm up (5 minutes)
- Static cycling (5minutes)
- Low resistance treadmill walking (5minutes)
- Play activities (5minutes)
- Cool down (5 minutes)

Proper rest periods were allotted between each sessions and was carried under the presence of the parent.

Relaxation technique group (*Annexure VI*)

The children in this group received Jacobson's relaxation technique which lasts for about 30 minutes a day for 3 alternate days a week for 4 weeks.

Jacobson's relaxation technique was demonstrated to the children in a quiet peaceful ambience. Then the children were asked to perform the same for about 30 minutes.

Training includes:

- Gentle breathing (5minutes)
- Visual imaging (5minutes)
- Tightening and loosening each body parts and joints (5minutes)
- Release state(5 minutes)
- Gentle breathing(5 minutes)
- Whole body stretching(5 minutes)

Proper rest periods were allotted between each sessions and was carried under the presence of the parent.

Materials

- Exercise mat
- Static cycle
- Treadmill
- Consent form
- Scoring chart
- Vitals monitoring instruments
- Stationary materials

5. DATA ANALYSIS& RESULTS

Data collected from participants of the same group (intra group) were analyzed using paired' test and the difference between the two groups (inter group) were analyzed using independent 't' test.

Paired't' test:

$$t = \frac{\bar{d}\sqrt{n}}{SD}$$

$$SD = \sqrt{\frac{\sum (d - \bar{d})^2}{n - 1}}$$

Where,

\bar{d} =calculated mean difference between pre and post-test values.

d =difference between pre and post-test values.

n =sample size

SD= standard deviation

Independent 't' test:

$$t = \frac{\bar{X}_1 - \bar{X}_2}{SD \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

$$SD = \sqrt{\frac{(n_1 - 1)SD_1^2 + (n_2 - 1)SD_2^2}{(n_1 + n_2) - 2}}$$

Where,

\bar{X}_1 = mean of group A

\bar{X}_2 = mean of group B

n_1 = number of subjects in group A

n_2 = number of subjects in group B

SD_1 = standard deviation of group A


SD_2 = standard deviation of group B

DATA INTERPRETATION

**Table I: Child Fatigue Scale (CFS) scoring for
Group A and Group B**

Groups		N	Mean	Mean difference	SD	The calculated 't' value
Group A	Pre	15	36.8	9.07	5.035	6.9686
	Post	15	27.73			
Group B	Pre	15	36.13	12.8	4.83	10.26
	Post	15	23.33			

**Graph I: Child Fatigue Scale (CFS) scoring for
Group A and Group B**



RESULTS:

GROUP A:

The mean and standard deviation of the group A is 9.07 and 5.035. The calculated 't' value is 6.9686 which is greater than the table value (1.76) at the level of $p < 0.05$. The result showed that there is significant difference between the pre and post-test of CFS score among participants of group A.


GROUP B:

The mean and standard deviation of group B is 12.8 and 4.83. The calculated 't' value is 10.26 which is greater than the table value (1.76) at the level of $p < 0.05$. The result showed that there is significant difference between the pre and post-test of CFS score among participants of group B.

**Table II: Parent Fatigue Scale (PFS) scoring for
Group A and Group B**

Groups		N	Mean	Mean difference	SD	The calculated 't' value
Group A	Pre	15	46	13.06	5.202	9.73
	Post	15	32.933			
Group B	Pre	15	42.267	13.6	6.378	10.69
	Post	15	28.667			

**Graph II: Parent Fatigue Scale (PFS) scoring for
Group A and Group B**



RESULTS:

GROUP A:

The mean and standard deviation of group A is 13.06 and 5.202. The calculated 't' value is 9.73 which is greater than the table value (1.76) at the level of $p < 0.05$. The result shows that there is a significant difference between the pre and post-test of PFS score among the participants of group A.


GROUP B:

The mean and standard deviation of group B is 13.6 and 6.738. The calculated 't' value is 10.69 which is greater than the table value (1.76) at the level of $p < 0.05$. The result shows that there is a significant difference between the pre and post-test of PFS score among the participants of group B

**Table III: Child Fatigue Scale scoring in
Aerobic vs. Relaxation group**

Parameters	Group	Mean	SD	Calculated 't' value	Table 't' value
Fatigue level	Group A	27.7	4.2770	2.848	2.048
	Group B	23.3			

**Graph III: Child Fatigue Scale scoring in Aerobic group vs.
Relaxation group**




RESULTS:

The mean values of aerobic training group (27.7) is greater than the relaxation training group (23.3). The calculated 't' value is 2.848 which is greater than the table value (2.048). The study showed that there is significant difference in child fatigue scale level of aerobic training group and relaxation training group at the level of $p < 0.05$. This shows that aerobic training is effective than the relaxation training in improvement of fatigue among ALL children.

**Table IV: Parent Fatigue Scale scoring in
Aerobic group vs. Relaxation group**

Parameters	Group	Mean	SD	Calculated 't' value	Table 't' value
Fatigue level	Group A	32.9	4.9940	2.3579	2.048
	Group B	28.6			

**Graph IV: Parent Fatigue Scale scoring in
Aerobic group vs. Relaxation group**



RESULTS:

The mean value of the aerobic training group (32.9) is greater than the relaxation group (28.6). The calculated 't' value is 2.3579 which is greater than the table value (2.048). The study showed that there is significant difference in parent fatigue scale level of aerobic training group and relaxation training group at the level of $p < 0.05$. This shows that aerobic training is effective than relaxation technique in improvement of fatigue in ALL children

6. DISCUSSION

The study was aimed to investigate the effects of Aerobic exercise and Relaxation technique on the children with Acute Lymphoblastic Leukemia (ALL). The ALL children have fatigue as one of the major problem which alters their mood and limits their functional activity. These children requires assistance for doing activities as they always seem tired, even with limited independent walking .When compared to normal child, they appear drained and gets exhausted easily. There are studies supporting the relationship between aerobic exercise and relaxation technique on fatigue level among children with ALL.

For the study purpose, 30 ALL children complaining fatigue were selected on the basis of inclusion criteria and divided into two equal groups. Both groups A and B were assigned for experimentation. Based on age parameter, 21 children were in the age group of 5-9 and 9 under the age group of 10-12. Based on the gender, 18 were boys and remaining 12 girls.

At the baseline of the study, all the participants were evaluated by the Child Fatigue Scale for measuring their level of fatigue. The parent fatigue scale questionnaire was filled by the parent and was

collected to measure the level of fatigue of their children. The collected data was taken as the pre interventional scores for analysis. Group A was treated with aerobic exercises. Group B was treated with Jacobson's relaxation technique. The treatment duration was over a period of 4 weeks. After this treatment period, all the participants were reevaluated by the same assessment tools and the collected data were taken as the post interventional scores for analysis.

The collected data was statistically analyzed in order to compare the variables, which was done within the groups by paired 't' test and to compare the variables between the groups which was done by the independent 't' test. The difference were considered at the significant level of $p < 0.05$.

The statistical analysis of variables within the group result showed significant difference. The **CSF** scoring result showed in Table 1 and Graph 1 accepted the alternative hypothesis 1 and 3 thus showing a significant difference in CFS outcome in aerobic and relaxation training groups among ALL children. The **PFS** scoring result showed in Table II and Graph II accepted the alternative hypothesis 2 and 4 thus showing a significant difference in PFS outcome in aerobic and relaxation training among ALL children.

The statistical analysis of the variables between aerobic exercise group and relaxation group result showed significant difference. **CSF** scoring result showed in table III and graph III accepted the alternative hypothesis 5 and confirmed the significant difference in outcome of CFS in aerobic and relaxation training group.

The PFS scoring result showed in table IV and Graph IV accepted the alternate hypothesis 6 and confirmed the significant difference in outcome of PFS in aerobic and relaxation training group.

The results confirmed the effects of Aerobic training which was effective than the relaxation technique in improving the cancer related fatigue. The result supported the previous studies which stated that aerobic training can improve fatigue in children. The study determined that there was significant increase in endurance level, which was also stated in the other studies. The improvement in fatigue level has a significant improvement in functional activities of daily living. All these factors contributed to the improvement in overall fitness, cardiovascular fitness, and increased endurance.

RECOMMENDATIONS

- Subjects with a large sample size can be studied.
- The effects on various age groups can be studied.
- Fatigue recovery of ALL children on various severity levels can be studied.
- Long term effects of aerobic training and relaxation training can be studied
- Tamil translated version of the parent and child fatigue questionnaire can be used to yield a better outcome as the original questionnaire is available in the English version.

7. CONCLUSION

The study confirmed the effects of both Aerobic exercise and Relaxation technique which could reduce the level of fatigue among children with ALL. The aerobic training yield better results compared to relaxation training in reducing fatigue level among ALL children.

REFERENCES

1. Frolayne. M. Carlos Wallace, Louping Zhang, Martyn .T. Smyth, Gabriella Rader and Craig Steinmaus. Parenteral, in utero and early life exposure to benzene and risk of Childhood Leukemia-A Meta-Analysis .April 27, 2015.
2. Pediatric physical therapy, 5th edition: Jan Stephen Tecklin, PT, MS. 2015, Lipincott Williams & Wilkins
3. Ghai Essential of Paediatrics,8th edition, OP ghai, 2013, CBS Publications
4. Victoria. M .Chia, Wilma M Schooner, Acute Lymphoblastic Leukemia: An assessment of international incidence, survival and disease burden. *Cancer cause control*, November 2015, issue 11, pp. 1627-1642.
5. Stancy .L Cooper; MD and Patrick A Brown, MD. Treatment of Pediatric Acute Lymphoblastic Leukemia .*Pediatric Clin NorthAm.*2015, Feb; 62(1):61-73.
6. Jennifer L Mcneer, Archie Bleyer, Valentino Conter, Wendy Stock. Cancer in Adolescents and young Adults: *Pediatric Oncology* pp. 151-175; Nov 18 2016.

7. Dhoriyani Narendra , Bhatt koushal D ,Smitha D ; A comparative study between Relaxation technique and Aerobic exercise in fatigue during chemotherapy in Acute Lymphoblastic Leukemia. *Indian Journal of Physiotherapy and Occupational Therapy* .July-September, Vol 7, no-3.
8. Efrat. L .Amitay PhD, MPH; Lital Keinan –Boker D.PhD Breastfeeding and childhood leukemia incidence-A Meta-Analysis and systemic review. *JAMA Pediatr.*2015; 169(6):e151025. Nov 2, 2015.
9. Karalexi M A, Dessypris, N Skalkidou A et al .Cancer causes control (2017)28:599
10. Robert B Gunier, Alice Kang, S Katherine Hammond, C Susanne Lea, Vonda Cruce, Jeffrey S Chang. A tasked based assessment of Parental occupational exposure to pesticides and childhood Acute Lymphoblastic Leukemia. *Environmental research*, vol 156, July 2017, pages 57-62.
11. Victoria G Margese, PhD, Lisa A Chiarello, PhD and Beverly J Lange. *Pediatric blood cancer*2004, 42:127-133.Effects of physical therapy intervention for children with acute lymphoblastic leukemia.

12. Dimeo F, Schwartz S, Fietz T, Wanjura T, Boning D, Theil E. Effects of endurance training on the physical performance of the patients with hematological malignancies during chemotherapy. *Journal of hematology and oncology, support carecancer* (2003)11:623.
13. Vogelzang NJ, Breitbart W, Cella D, Curt G ,Groopman J E. Fatigue in cancer, *BMJ*, 2001;322;1560. Vol 322 30 June 2001.
14. Mock V, Dow K H, Meares C J, Grimm P M, Dienemann J A, Haisfield Wolfe ME, Quitasol W and Mitchell S. Effects of exercise on fatigue, physical functioning and emotional distress during radiation therapy for breast cancer. July 1997, 24(6):991/1000.
15. Anna Spathis, MSc, Sara Booth, Sarah Grove, M B Chir, Isla Kuhn, MSc, and Stephen Barclay M .Cancer related fatigue practice and future research .University of Cambridge, United Kingdom.
16. Karen Seiter, MD; Emmanuel C Besa, MD. Acute Lymphoblastic Leukemia medication .Updated June 23, 2017.

17. Ching –hon Pui ,MD and William E Evans, Pharm D. Drug therapy-treatment of Acute Lymphoblastic Leukemia. *The NewEngland Journal of Medicine*, N Engl J Med 2006; 354: 166-178.
18. Jabeen Kishwer MBBS, Ashraf Mohammed S MD, MBBS, DCH, MRCPI; Ifthikar, Sundus MS, Belgaumi Asim F, MD. The impact of socioeconomic status on the outcome of childhood ALL, treatment in low/middle income country. *Journal of pediatric hematology/oncology*, Nov 2016- volume 38 issue 8-p 587-596.
19. Vanesa Caruso, Licia Lacoviello, Augusto Di Castelnuovo et al. Thrombotic complications in childhood Acute Lymphoblastic Leukemia: A Meta-Analysis of 17 prospective studies comprising 1752 pediatric patients. *Blood* 2006 108:2216-222;
20. Lidewji T Warris, Marry M van den Heuvel Eibrink, Femke K Arsen Cor van den Bos, Christian M Swaan, Rob Pieters and Erica L T van den Akker. A double blind randomized control trial to prevent serious side effects of Dexamethasone during patient treatment of ALL .*Blood* 2015, vol \126:2495 issue 23.

21. Leonardo A Mattano, Harland N Sather, Michael E Trigg, James B Nachman. *Journal of clinical oncology*; Osteonecrosis as a complication of treating Acute Lymphoblastic Leukemia in children: A report from the children cancer group. .
22. Lip Shultz S E, Colan S D, Gelber R D, Perez-Atayde. A .R, Sallan S E, Sanders S P .Late cardiac effects of Acute Lymphoblastic Leukemia in childhood. Eng. JMed.1991 Mar21; 324(12):808-15.
23. Yin Ting Cheung, Kevin R Krull .Neurocognitive outcomes in long term survivors of child Acute Lymphoblastic Leukemia treated on contemporary treatment protocols: A systemic review. j.neubiorev.2015.03.016
24. Marja Schoenmakers, Tim Takken, Vincent A M Gulmans, Nico LU Van Meetren et al. Muscle strength and functional ability in children during and after treatment for acute lymphoblastic leukemia or T cell Hodgkin lymphoma- a pilot study .*Department of pediatric physical therapy and exercise physiology*. July 31, 2006.
25. James V Tricoli, Donald G Blair, Cardey K Anders .Biological and clinical characteristics of adolescent and young adult

- survivors. : Acute Lymphoblastic Leukemia, Colorectal cancer, Breast Cancer, Melanoma and Sarcoma. American Cancer Society, 2016; 122:1078-28.
26. Shadi Farzin Gohar MD, Melanie Comito M D, Jennifer Price and Victoria Marchese PT, PhD .Feasibility and parent satisfaction of a physical therapy intervention program for children with Acute Lymphoblastic Leukemia in the first 6 months of medical treatment. *Pediatric blood and cancer*, volume 56, issue 5 may 2011; pp 799-804.
 27. Michele ,Sujan Hong ,Cheryl L Cox, Wendy M Leisenring, Kevin C Oeffinger ,Jill Ginsberg ,Gregory T Amstrong ,Kirstein k ness ,Leslie L Robinson. Physical therapy and chiropractic use among childhood cancer survivors with chronic disease: Impact on health related quality of life.
 28. Victoria G Marghese PhD, Lisa A Chiarello, Beverly J Lange M D. Effects of physical therapy intervention for children with Acute Lymphoblastic Leukemia. *Pediatric blood and cancer*, volume 42, issue: 2 Feb 2004; pp 127-133.
 29. Spelman, Andrea Buck SPT; Harris, Jessica SPT. Effects of exercise on quality of life and function in patients with

Leukemia currently undergoing treatment: a Systemic Review of Literature. *Rehabilitation Oncology*: 2015-volume 33 issue 3, p 6-18.

30. Maria Beatriz Perondi, Bruno Gualano, Vitor de Salles. Effect of combined aerobic and strength training program in youth patients with Acute Lymphoblastic Leukemia. *Journal of sportscience and medicine*. Sports sci Med.2012 Sep; 11(3):387-392.
31. Dimeo F, Fetscher S, Lange W, Mertelsmann R, Keul J. Effects of Aerobic exercise on the physical performance and incidence of treatment related complications after high dose chemotherapy. *Journal of hematology and oncology*. 1997, volume 90:3390-3394.
32. Julienne E Bower, Kate Bak, Ann Berger, William Breitbart. Screening assessment and management of fatigue in adult survivors of cancer: An American Society of Clinical Oncology clinical practice guideline adaptation. *Journal of clinical oncology, ASCO articles*. No 17(June 2014) 1840-1850.
33. Marilyn J Hockenberry, Pamela S Hinds, Patrick Barrera, BS, Rosalind Bryant, MSN, PNP. Three instruments to assess

fatigue in children with Cancer: The Child, Parent and Staff perspectives ;*Journal of pain and symptom management* , Vol 25,No 4 April 2003.

34. Measures of fatigue and sleep; Geri Neuberger. *American College of Rheumatology*, Vol 49, no 55, Oct 15 2003,pp S175-S183
35. San Juan, Alejandra F, Fleck, Steven J, Chamorro –vina, Carolina; Claudia; Mari Fernandez. Effects of an intra-hospital exercise program intervention for children with leukemia. April 2006.
36. Tseng-Tieng Haung and Kirsten K Ness. *Department of Epidemiology and Cancer control*. Exercise intervention in children with cancer: A review; September, 3, 2011.
37. Fernando C Dimeo, Frank Thomas, Felix Propper, Michael Mathias, Cornelia Raabe –Menssen. Effects of Aerobic exercise and Relaxation training on fatigue and physical performance of cancer patients after surgery. A randomized control trial. *Support cancer care* (2004).volume 12, issue 11, pp774-779.

38. C H Pui, D Pei, D Campana, C Cheng, J T Sandlund, W P Bowman. A revised definition for cure of childhood Acute Lymphoblastic Leukemia. *Leukemia* (30 April 2014)
39. Deborah Tomlinson ,MN ,RN ,Pamela S Hinds, Marie – Chantal Either ,Kirstein k Ness PT ,PhD, Sue Zupanec .Psychometric properties of instruments used to measure fatigue in children and adolescent with cancer: a systemic review. *Journal of pain and symptom management*, Vol 45 no 1, January 2013.
40. Ladha ,Aliya B MSc; Courneya , Kerry S PhD ;Bell ,Gordon J Field, Catherine J PhD ;Grundy,Paul M D .Effects of acute exercise on neutrophils in pediatric Acute Lymphoblastic Leukemia survivors.2006 Lippincott Williams &Wilkins,Inc.
41. Dimeo, Fernando, Rumberger, Brigitta G, Keul Josh .*Medicine science in sports &exercise* .vol 30 (4) April 1998, 475-478 .Aerobic exercise as a therapy for cancer fatigue.
42. Moyer Mileur, Laurie J, PhD, RD; Randsell, Lynda PhD, Bruggers, Carol S MD. Fitness of children with standard risk of Acute Lymphoblastic Leukemia during maintenance therapy.

43. Relaxation techniques; a practical handbook for health care professional. Rosemary A Payne Churchill Livingstone, 1998.
44. Pediatric essential physiology: Neil, Spurway, and Don Maclaren Elsevier publications.
45. ACSMS (American College of sports medicine) guidelines for exercises resting and prescription; 7th edition. Lawrence Amstrong ,PhD ;Gary J Balady M D; Michael J Berry ,PhD.S
46. Ali Hassanpour-Dehkordi and Amir Jalali. Effect of progressive muscle relaxation on the fatigue and QOL among Iranian aging persons. *Acta Medica Iranica*, 2016, 54(7):430-436.
47. Alison Crichton, Sarah knight, Ed Oakley, Franz E Badi, M D. fatigue in children chronic health conditions: a systematic review of assessment instruments *.Journal of pediatrics*, vol 135, no 4, April 2015
- 48.** Donald P Lawrene , Joseph Lan, Kimberly Miller Evidence Report on the Occurrence, Assessment, and Treatment of Fatigue in Cancer Patients *Journal of National Cancer Institute Monographs* ,no 32, 2004.

ANNEXURE-I
CONSENT FORM

I..... (parent) hereby agree to provide my fullest consent and co-operation to allow my child to be taken as a subject for the research work of (Reg no.271540062) entitled “*The efficacy of aerobic exercises and relaxation technique in fatigue among children with acute lymphoblastic leukemia-A comparative study*”. I have decided to volunteer my child for the study on my own will and was not compelled by individual or group of people and my consent is not for any monetary benefits.

The possible outcomes and effects of the study as well as the procedure that will be executed on my child is fully explained to me by an investigator in the language best known to me and I am aware that my child being subjected to this study and I’ll have to give more time for assessments and treatments.

The questions and queries I have posed have been answered to my satisfaction and I am aware that my child’s identity will be kept confidential. I am also aware that I can discontinue or withdraw from this study on my child at anytime without adversely affecting my child’s health.

The matter in this consent form was read by me /read to me by an investigator and is true and understanding to the fullest of my knowledge.

Sign of the parent

Sign of the investigator

Place:

Date :

ANNEXURE -II
ASSESSMENT FORM

Name:

GROUP: ID NO:

Age:

Date of assessment:

Gender:

Height:

Weight:

Informant:

Medical history:

Chief complaints:

Present medical illness:

Onset:

Duration of hospital stay:

Birth history:

Developmental history:

Immunization history:

Drug history:

Higher mental functions:

Cognitive development:

Socially adaptive:

Speech:

Vision:

Hearing:

Vitals:

Temperature:

Pulse rate:

SPO2:

BP:

RR:

General appearance:

Facial:

Posture:

Behavior:

Sleep activity:

Appetite:

Interactiveness:

General health status:

Breathing:

Muscle tone:

Circulation:

Nutritional status:

Hydration:

Mental state:

Level of fatigue:

Hematological:

Platelet count:

WBC count:

Units of blood transfused:

Neurological:

Level of consciousness:

Gait:

Recent changes:

Changes in appearance:

Changes in treatment:

ANNEXURE –III
CHILD FATIGUE SCALE

Sl No	How have you been feeling during the past one week?	Yes/no	Not at all	A little	Someti mes	Quite a lot	A lot
1.	I have been tired.						
2	My body has felt different.						
3	I have been tired in the morning.						
4	I need a nap most of the time						
5	I have not been able to play.						
6	I have been playing around.						
7	I have been feeling sad.						
8	I have been mad.						
9	I stop doing things and take rest.						
10	I have not been able to do activities.						
11	I have not been able to run.						
12	I found hard to keep my eyes open.						
13	I have slept more at night						
14	I have trouble thinking.						
Score: 1-not at all 2-a little 3-some times 4-quite a lot 5-a lot							

TOTAL SCORE:

INTERPRETATION

The *Childhood Fatigue Scale (CFS)* is a 14-item, two-part instrument. The items ask the child for a “yes” or “no” (frequency) response regarding their experience of any fatigue-related symptoms during the past week. If the statement is true for the child, he or she is asked to rate how much the problem bothers the child on a five-point Likert scale ranging from “Not at all” to “A lot.” (Intensity). If the child has not experienced the particular problem, and has answered with a “no” response for the question, the score is zero for that question. Frequency scores range from 0 to 14, and intensity scores, also considered total fatigue scores, ranges from 0 to 70. Higher scores correspond to greater amounts of experienced fatigue.

ANNEXURE –IV
PARENT FATIGUE SCALE

Sl.No	How your child have been feeling during the past week?	Not at all	Almost never	Sometimes	Not always	Always
1.	My child has been tired in the morning.					
2.	My child has hard time getting out of bed.					
3.	My child has been too tired to eat.					
4.	My child has not slept throughout night.					
5.	My child has been tired in the afternoon.					
6.	My child always needed a nap.					
7.	My child seems to have no energy.					
8.	My child feels tired after night's sleep.					
9.	My child has play changes during the day.					
10.	My child always wanted to lie down.					
11.	My child needs rest while walking.					
12.	My child has become quieter.					
13.	My child has become less interactive.					
14.	My child has become more irritable.					
15.	My child shows mood swings.					
16.	My child has become in co-operative.					
17.	My child has dark circles under his/her eyes.					
Score:		1-not at all	2-almost never	3-sometimes		
		4-not always	5-always			

TOTAL SCORE:

INTERPRETATION

The *Parent Fatigue Scale (PFS)* consists of 17 items regarding parents' perceptions of the amount of fatigue experienced by their children in the past week. The items are rated on a five point Likert scale ranging from 1-“Not at all “to 5-“Always” . Scores on the parent scale range from 17–85 with higher score corresponding to greater amounts of perceived fatigue.

ANNEXURE –V
TREATMENT PROTOCOL FOR AEROBIC
EXERCISE GROUP

Aerobic training may be defined as the ability to deliver oxygen to the exercising muscles and utilize it to generate energy during exercise. Aerobic fitness depends on pulmonary, cardiovascular and hematological components of oxygen delivery and oxidative mechanism of exercising muscle.⁽⁴⁴⁾

GUIDELINES:

- Exercises should be performed at least after 2hrs after consumption of food.
- Should not have exercised vigorously prior to the treatment session.
- Should be made familiar with treadmill and static cycling.
- Procedures should be well explained to the children.
- The vitals should be checked prior to exercise sessions.
- The safety and well-being of the children are of great importance as they are immune- compromised.

Treatment sessions:

1. Initial warm up

Duration: 5 minutes

Low intensity warm ups are advisable.

These includes: speed walking and arm swinging

Initial starting position: Standing, with arms on sides.

Instructions: Ask the children to start walking with their maximum speed with arm swinging on sides from their starting position.

Trainer: Stands and observes the children; counts on stop watch. After 5 mints general warm-up, provide rest for about 2 minutes.

2. Static cycling

Duration: 5 minutes

Initial starting position: Sitting upright, heels on the pedal, knees fully extended, arms on grab bars.

Instructions: Ask the child to pedal up the cycle slowly and gradually increase the speed.

Trainer: Stand nearby the child and set the timer. After 5 minutes of cycling, provide rest for about 2 minutes.

3. Low resistance treadmill walking Duration: 5 minutes

Initial starting position: Stand upright, hands on grab bars, feet flat on the platform.

Instructions: Ask the child to start walking slowly with heads upright.

Resistance should be appropriate to the size and age of the children.

Trainer: Stand near the child, monitor the activity and set the timer.

Monitor Heart Rate (HR), maximum of about 195. After treadmill walking for 5 minutes, provide proper rest period for about 2 minutes.

4. Play activities duration: 5 minutes

Children are made to indulge in group activities to bring relief and to make them overcome the sense of training. Activities like jogging, passing ball, chasing, simple aerobic dance steps are performed in groups (any of these).

Initial starting position: Standing on pairs (groups of 2)

Instructions: Ask the children to start jogging slowly standing face to face.

Trainer: Stand near the groups and observe the activities. Set the timer.

5. Cool down

Duration: 5 minutes

Cool down sessions should be made more interactive and relaxing. Activities of gentle breathing and stretching are incorporated. Children are asked to slow down jogging into mild static walking to lower their breathing rate and gradually come to pause. Children are made to take 2/3 sets of breathing.

Initial starting position: standing with arms on side.

Instructions: Gently breathe in and bring both the hands outward and over the head. Gently breathe out and bring back the hands to the sides of the body. Make 3-5 repetitions.

Trainer: stand near the children and observe their activities.

Benefits of aerobic exercises:

- Improves general fitness.
- Tones muscle groups.
- Increases bone strength.
- Increases cardiovascular fitness.
- Increases oxygen carrying capacity.
- Reduce stress and depression.

These techniques can be easily performed by the children and are easy to administer. When incorporated with play activities and in groups, children get more interested in performing it.

ANNEXURE-VI
TREATMENT PROTOCOL FOR RELAXATION TRAINING
JACBSONS PROGRESSIVE RELAXATION

This technique initially found by Edmund Jacobson (1938) and later modified by Wolpe (1958) and later Bernstein and Borkoveks (1973) was named as ‘progressive relaxation training’ (PRT).⁽⁴³⁾

PRT is defined as learning to relax specific muscle groups while paying attention to the feelings associated with both the tensed and relaxed states. The aims are:

- To achieve a state of deep relaxation in increasingly short periods.
- To control excess stress in stress inducing conditions.

Jacobson’s progressive relaxation technique involves contracting and relaxing the body parts to feel calmer. Before starting the sessions, a general format of this technique should be well explained to the children. Initially it might be difficult to relax the parts but with proper training and time, they can master these technique and can practice throughout life.

PROCEDURE:

- A silent and calm environment is necessary to perform training.
- The hospital setting with limited crowd area is selected for the same.
- The children are made familiar with the method, explained and demonstrated well before the actual starting of the sessions.
- The children are made to lie down freely in the mat and are advised to wear loose clothes. This takes around 5 mints.

First session:

duration: 5 minutes

1. Gentle breathing

- Children are advised to close their eyes.
- Ask the children to observe the natural rhythm of breathing.
- Ask them to take a full breath of air in -hold –and let go. Repeat 3-5 times.
- Inhale through the nose, raise your tummy and slowly exhale through the mouth.

Second session:

Duration: 5 minutes

2. Visual imaging

- Ask the children to visualize their body parts
- Starting from toes, knees, thighs, buttocks, stomach, chest, shoulder, neck, hands, face and head.
- Ask them to visualize objects like birds, trees, school, home, waterfalls and so on.
- Ask them to listen the sounds like twitching of birds or tapping of foot.

Third session

Duration: 5 minutes

3. Tightening and loosening of body parts

- Guide the children with proper instructions for each body part:
- Gently pull up the toes towards body-tighten-hold up-loosen.
- Press heels on the floor-tighten-holdup-loosen.
- Pull knees together –hold briefly-loosen.
- Squeeze the buttocks together-hold –loosen.
- Gently pull in the tummy muscles –tighten –holdup-loosen
- Gently pull the shoulders up tight towards the ears-hold-loosen.
- Gently press the elbows and upper arms to the side of the body-hold –loosen.
- Hands gently clench, tighten-hold-loosen.

- Push the head forward slightly –hold up-let the head go back to normal position.
- Grit the teeth together-hold-loosen jaw.
- Purse the lips –hold up tightly-loosen.
- Press the tongue to the roof of the mouth –hold-loosen
- Tighten the eyes-hold-loosen.
- Forehead frown a little-hold-loosen.

Forth session

Duration: 5 minutes

4. Release state

- This session is more free and enjoying.
- Ask the children to feel the difference from the previous session
- Ask them to spend a few moments enjoying the feeling of release, releasing a little more with each exhale.

Fifth session

Duration: 5minutes

5. Gentle breathing

- Ask the children to become aware of the present breathing pattern, contact between the body and floor/bed.
- Ask the children to gently open the eyes.
- Ask them whether they feel any sort of bodily changes after the session.

Sixth session

duration: 5 minutes

6. Stretching

- Ask the children to gently bring the palms together, extend and bring back of their head.
- Ask them to pull their hands upward and push their legs downward and feel the whole body stretching. Hold for a few seconds and release.
- Ask the children to get up and have their activities.